

# Northumbria Research Link

Citation: Immerzeel, W. W., Lutz, A. F., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S., Davies, B. J., Elmore, A. C., Emmer, A., Feng, M., Fernández, A., Haritashya, U., Kargel, J. S., Koppes, M., Kraaijenbrink, P. D. A., Kulkarni, A. V., Mayewski, P. A., Nepal, S., Pacheco, P., Painter, T. H., Pellicciotti, Francesca, Rajaram, H., Rupper, S., Sinisalo, A., Shrestha, A. B., Viviroli, D., Wada, Y., Xiao, C., Yao, T. and Baillie, J. E. M. (2020) Importance and vulnerability of the world's water towers. *Nature*, 577 (7790). pp. 364-369. ISSN 0028-0836

Published by: Nature Publishing

URL: <https://doi.org/10.1038/s41586-019-1822-y> <<https://doi.org/10.1038/s41586-019-1822-y>>

This version was downloaded from Northumbria Research Link:  
<http://nrl.northumbria.ac.uk/id/eprint/44492/>

Northumbria University has developed Northumbria Research Link (NRL) to enable users to access the University's research output. Copyright © and moral rights for items on NRL are retained by the individual author(s) and/or other copyright owners. Single copies of full items can be reproduced, displayed or performed, and given to third parties in any format or medium for personal research or study, educational, or not-for-profit purposes without prior permission or charge, provided the authors, title and full bibliographic details are given, as well as a hyperlink and/or URL to the original metadata page. The content must not be changed in any way. Full items must not be sold commercially in any format or medium without formal permission of the copyright holder. The full policy is available online: <http://nrl.northumbria.ac.uk/policies.html>

This document may differ from the final, published version of the research and has been made available online in accordance with publisher policies. To read and/or cite from the published version of the research, please visit the publisher's website (a subscription may be required.)



**Northumbria  
University**  
NEWCASTLE



University**Library**

# Importance and vulnerability of the world's water towers

*Immerzeel, W.W., Lutz, A.F., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S.,  
Davies, B.J., Elmore, A.C., Emmer, A., Feng, M., Fernández, A., Haritashya, U., Kargel, J.S., Koppes, M.,  
Kraaijenbrink, P.D.A., Kulkarni, A.V., Mayewski, P., Nepal, S., Pacheco, P., Painter, T.H., Pellicciotti, F.,  
Rajaram, H., Rupper, S., Sinisalo, A., Shrestha, A.B., Viviroli, D., Wada, Y., Xiao, C., Yao, T., Baillie,  
J.E.M.*

Mountains are the water towers of the world, sustaining a substantial part of both natural and anthropogenic water demands. These water towers are highly sensitive and prone to climate change, yet their importance and vulnerability have not been quantified at the global scale. Here, we develop and present the first global Water Tower Index, which ranks all water towers in terms of their water supplying role and the downstream dependence of ecosystems and society. For each of them, we assess its vulnerability related to water stress, governance, hydropolitical tension, and future climatic and socio-economic changes. We conclude that the most important water towers, predominantly located in Asia, are also among the most vulnerable, and that climatic and socio-economic changes will impact them profoundly. This could negatively impact 1.9 billion people living in (0.3 billion) or directly downstream (1.6 billion) of mountains. Immediate action is required to safeguard the future of the most important and vulnerable water towers of the world.


## 21    **Global Water Towers**

22    The term “water tower” is used to describe the role of water storage and supply that mountain  
23    ranges play to sustain environmental and human water demands downstream<sup>1,2</sup>. Compared to its  
24    downstream area, a water tower (seasonally) generates higher runoff from rain as a result of  
25    orographic precipitation and delays the release of water by storing it in snow and glaciers (because of  
26    lower temperatures at high altitude) and lake reserves. Because of their buffering capacity, for  
27    instance by supplying glacier melt water during the hot and dry season, water towers provide a  
28    relatively constant water supply to downstream areas. We define a water tower unit (WTU; see  
29    Methods, Extended Data Figure 1) as the intersection between major river basins<sup>3</sup> and a topographic  
30    mountain classification based on elevation and surface roughness<sup>4</sup>. Since water supply and demand  
31    are linked at the river basin scale, the basin is the basis for the WTU. One WTU can therefore contain  
32    multiple topographically different mountain ranges and we assume that it provides water to the  
33    areas in the downstream river basin that are hydrologically connected to the WTU (Extended Data  
34    Figure 1, Extended Data Table 1 and 2). Subsequently, we only consider cryospheric WTUs by  
35    imposing thresholds on satellite derived snow cover data<sup>5</sup> and a glacier inventory<sup>6</sup>, because the  
36    buffering role of glaciers and snow and the delayed supply of melt water is a defining feature of  
37    water towers. Consequently, there are regions (e.g. in Africa), which do contain mountain ranges, but  
38    because of their small snow and ice reserves they do not meet the WTU criteria. In total, we define  
39    78 WTUs globally (see Methods), which are home to more than 250 million people. However, more  
40    than 1.6 billion people live in areas receiving water from WTUs, which is about 22% of the global  
41    population<sup>7</sup> (Figure 1).

42    Water towers play an essential role in the Earth system and are particularly important in the global  
43    water cycle<sup>1,2</sup>. In addition to their water supply role, they provide a range of other services<sup>8,9</sup>. About  
44    50% of the global biodiversity hotspots on the planet are located in mountain regions<sup>10</sup>, they contain  
45    a third of the entire terrestrial species diversity<sup>11</sup>, and are extraordinarily rich in plant diversity<sup>12</sup>.  
46    Moreover, mountain ecosystems provide key resources for human livelihoods, host important



47 cultural and religious sites, and attract millions of tourists globally<sup>4</sup>. Economically, 4% and 18% of the  
 48 global Gross Domestic Product (GDP) is generated in WTUs and WTU-dependent basins  
 49 respectively<sup>13</sup>. Furthermore, mountains are highly sensitive to climate change<sup>14,15</sup> and are warming  
 50 faster than low-lying areas due to elevation-dependent warming<sup>16</sup>. Climate change therefore  
 51 threatens the entire mountain ecosystem. Worldwide, the vast majority of glaciers are losing mass<sup>17</sup>,  
 52 snow melt dynamics are being perturbed<sup>18–21</sup>, and precipitation and evapotranspiration patterns are  
 53 shifting, all leading to future changes in the timing and magnitude of mountain water availability<sup>22</sup>.



54  
 55 **Figure 1: The Water Tower Index (WTI), population in WTUs and their downstream basins. The WTI, derived from the**  
 56 **supply and demand index, is shown for all 78 Water Tower Units (WTU), in combination with the shaded total population**  
 57 **in all WTU-dependent river basins. Labels indicate the five water towers with the highest WTI value per continent. The**  
 58 **insets show the number of people living in WTUs as a function of elevation and the downstream population's proximity**  
 59 **to the WTUs<sup>7</sup>.**

60 Not only are the world's water towers crucial to human and ecosystem survival, the steep terrain in  
 61 combination with extreme climatic conditions, and in some regions seismic or volcanic activity,  
 62 frequently triggers landslides, rock fall, debris flows, avalanches, glacier hazards and floods<sup>23,24</sup>. Since  
 63 2000 alone, over 200,000 people have died in WTUs as a result of natural disasters<sup>25</sup>. Climate change,  
 64 in combination with population growth, urbanization, and economic and infrastructural

65 developments is likely to exacerbate the impact of natural hazards and further increase the  
66 vulnerability of these water towers<sup>26–30</sup>.

## 67 **Quantifying importance of water towers**

68 Consequently, there is a strong need for a consistent framework to assess and rank the importance  
69 and vulnerability of individual WTUs in order to guide global research, as well as conservation and  
70 policy-making efforts. Here we develop such a framework according to quantifiable indicators for  
71 both the water supply and demand sides of each WTU. Conceptually, a WTU is deemed important  
72 when its water resources (liquid or frozen) are plentiful relative to its downstream water availability  
73 and when its basin water demand is high and cannot be met by downstream water availability alone.  
74 Ideally, such an assessment would require a global-scale, high-resolution, fully coupled atmospheric-  
75 cryospheric-hydrological model that can resolve the interactions between extreme topography and  
76 the atmosphere, fully account for snow and ice dynamics, and incorporate anthropogenic  
77 interventions in the hydrological cycle. It would also require models that include socio-economic  
78 impacts on sectoral water demands and a spatially explicit attribution of water source (e.g.  
79 meltwater, groundwater, surface runoff) to water use. Although important progress has been made  
80 in specific regions and for specific sectors<sup>31</sup>, at global scale this is not yet feasible. We therefore  
81 derive indices covering relevant drivers for both the water supply and demand of a WTU's water  
82 budget (see Methods), which we combine to derive a Water Tower Index (WTI).

83 The supply index (SI) is based on the average of four indicators that are quantified for each WTU:  
84 precipitation, snow cover, glacier ice storage and surface water (Figure 2a, Extended Data Table 3,  
85 Table S1, Methods). If the precipitation in the WTU (Extended Data Figure 3a) is high relative to the  
86 overall basin precipitation and if the inter-annual and intra-annual variation is low (i.e. the supply is  
87 constant), a WTU scores high on the precipitation indicator. If a WTU has persistent snow cover  
88 (Extended Data Figure 3b) throughout the year and the snowpack shows lower inter-annual  
89 variation, this will result in a high snow indicator. Similarly, if the total glacier ice volume (Extended

90 Data Figure 4a) and glacier melt water yield in a WTU are high relative to the basin precipitation then  
91 a WTU has a high glacier indicator value. Finally, we assess the amount of water stored in lakes and  
92 reservoirs in a WTU (Extended Data Figure 4b) compared to basin precipitation to derive a surface  
93 water indicator.

94 There is considerable variability in the power of WTUs to supply water. In Asia, the Tibetan Plateau  
95 ranks highest because of the large amounts of water stored in lakes, but a large part of the Tibetan  
96 Plateau is endorheic and its water resources are disconnected from the downstream demand. The  
97 Indus WTU has an important water supplying role with a balanced mix in the importance of  
98 precipitation, glaciers, snow and surface water. In Europe, the Arctic Ocean Islands, Iceland, and  
99 Scandinavia have extensive stocks of water stored in their WTUs. Iceland stands out with some of the  
100 thickest glaciers in the world and a glacier ice storage ( $\sim 1,027 \text{ km}^3$ ) that is 15 times as large as its  
101 total annual WTU precipitation ( $\sim 67 \text{ km}^3$ ). In South America, the mountain ranges (Extended Data  
102 Table 1 and 2) supplying the the Southern Chilean Pacific coast regions and La Puna Region are the  
103 most prominent water towers, because of large glacier ice reserves and high orographic precipitation  
104 rates and due to large amount of water stored in lakes in the case of La Puna region. The Northwest  
105 Territories and Nunavut, Fraser, and the Pacific and Arctic coast are the key WTUs in North America.  
106 In the Northwest Territories and Nunavut the significance of the WTU is primarily driven by the  
107 abundance of glaciers, snow, and surface water. However, the precipitation indicator value is low,  
108 meaning that mountain precipitation is low relative to the overall basin precipitation.





Figure 2: The supply and demand index. The supply index (SI; panel a) and demand index (DI; panel b) of each WTU grouped by continent and ordered by SI or DI value, respectively. Increasing radially, the stacked bars show the four indicator values for surface water (L), glacier (G), snow (S) and precipitation (P), respectively. In panel b, increasing radially, the stacked bars show the four indicator values for natural ( $D_{NAT}$ ), industrial ( $D_{IND}$ ), domestic ( $D_{DOM}$ ) and irrigation demands ( $D_{IRR}$ ), respectively. Calculation details of the indicators and index of the indices are provided in Extended Data Table 3 and Extended Data Table 4.

To derive a demand index (DI) for each WTU, we quantify the monthly water requirements to be supplied by the water towers to sustain the WTU basin's net sectoral water demand for irrigation, industrial (energy and manufacturing) and domestic purposes, and monthly natural water demand, relative to the total annual demand (Figure 2b, Extended Data Table 4, Table S1). Monthly sectoral water requirements are estimated by subtracting the monthly water availability downstream (ERA5 precipitation minus natural evapotranspiration ( $P-ET$ )<sup>32</sup>) from the monthly net demands<sup>33</sup>. The demand index is the average of the four indicators (see Methods). Figure 2b demonstrates considerable variability, globally and within continents, in the demands that WTUs need to sustain. Irrigation water demands are the highest of the four demand types, and this is relatively consistent across the continents. The Asian river basins, specifically the heavily irrigated and densely populated basins such as the Indus, Amu Darya, Tigris, Ganges-Brahmaputra, and Tarim, score higher on the demand index than other basins across the world and they score high on each sectoral demand indicator. In those basins, the required water to close the gap between demand and downstream supply may also originate from (unsustainable) groundwater use<sup>34,35</sup>. However, specifically in those cases, when there is a large water gap being (partly) closed by unsustainable groundwater pumping, the WTU water supply is critical to both meet the demand and recharge the aquifers.

In Europe, the Volga and Ural in Russia show the highest demand index values, including high values for the natural demand indicator, whereas the Negro basin stands out in South America. In North America a range of basins scores equally high, but for different reasons. For example, the Mississippi-

Missouri basin scores high particularly because of a high natural demand indicator value, whereas the California basin scores high on all four demand indicators.

Ultimately, the presence of mountain water resources, either as additional rain or stored in snow, ice, or lakes, in conjunction with a high demand downstream, determines whether a WTU has an indispensable role (Extended Data Figure 2). The Water Tower Index (WTI) is the product of SI and DI, for which the values are subsequently normalized over the range of WTI values found for all 78 WTUs (Figure 1, Table S1). Globally, the upper Indus basin is the most critical water tower unit (WTI=1.00±0.03) with abundant water resources in the Karakoram, Hindu-Kush, Ladakh, and Himalayan mountain ranges in combination with a densely populated and intensively irrigated downstream basin<sup>22,36</sup>. In North America, the Fraser and Columbia River Basins are the most critical WTUs (WTI=0.62±0.07 and 0.58±0.06, respectively). The Fraser Basin is rich in surface water resources, and has a high natural water demand downstream, whereas the Columbia Basin is rich in snow and glacier resources in combination with a high irrigation demand. In South America, the Cordillera Principal, the Cordillera Patagónica Sur, and the Patagonian Andes are key WTUs in the supply of water to the South Atlantic and Pacific Coasts regions and the Negro basin. In Europe, the Alps are the most relevant water supplying mountain range to meet the demands of the Rhône, (WTI=0.45±0.07), Po (WTI=0.39±0.07) and Rhine (WTI=0.32±0.11) basins. Note that several WTUs that score high on either the supply index or the demand index do not rank high in the final Water Tower Index. For example, the Tibetan Plateau and Arctic Ocean Islands WTUs score high on the supply index, but have the lowest scores on the demand index, due to low water demands (Figure 2b). By contrast, the Sabarmati in Asia with a small portion of its water coming from the Himalayas scores highest on the demand index, but low on the supply index.


## **Vulnerability of the water towers**

We assess the vulnerability of each WTU and show this for the five most important (i.e. with highest WTI values) WTUs in Asia & Oceania, Europe, North America, and South America (Figure 3, Table S2).

For this analysis, we include the hydro-political tension<sup>37</sup>, baseline water stress<sup>38</sup>, government effectiveness<sup>39</sup>, projected climate change<sup>40</sup>, projected change in GDP<sup>41</sup>, and projected population change<sup>7</sup> (see Methods). The highest ranking WTUs of South America and Asia in particular are more vulnerable than those in North America and Europe. Strikingly, the Indus, which is globally the most important water tower (Figure 4), is also very vulnerable. The Indus is a transboundary basin with considerable hydro-political tension between its riparian countries Pakistan, India, China, and Afghanistan. The population of ~206 million people in the basin in 2016 is projected to increase by 50% until 2050, and the basin's GDP is projected to encounter a nearly eightfold increase<sup>41</sup>. The temperature in the Indus WTU is projected to increase by 1.9 °C between 2000 and 2050, compared to 1.8 °C in the downstream section<sup>40</sup>. The average annual precipitation in the Indus WTU is projected to increase by 0.2%, compared to 1.4% downstream<sup>40</sup>. It is evident that, due to the expected strong growth in population and economic development, the demand for fresh water will rise exponentially<sup>42</sup>. Combined with increased climate change pressure on the Indus headwaters, an already high baseline water stress and limited government effectiveness, it is uncertain whether the basin can fulfil its water tower role within its environmental boundaries. It is unlikely that the Indus WTU can sustain this pressure.

The Indus does not stand alone, however. Nearly all important WTUs in Asia are also highly vulnerable (Figure 3). Most WTUs are transboundary, densely populated, heavily irrigated basins and the vulnerability is primarily driven by high population and economic growth rates and in most cases ineffective governance. Moreover, the Syr Darya, Amu Darya and Indus, in particular, are characterized by considerable hydro-political tension<sup>37</sup>. In most cases, downstream riparian states are dependent on mountain water resources provided by bordering upstream states to supply the competing irrigation, hydropower and domestic demands. In South America, the vulnerability is less than for the Asian WTUs, and the drivers are variable. In North Chile Pacific Coast, the baseline water stress and a projected decrease in precipitation (-4.8%) cause the vulnerability, whereas in La Puna

184 Region population and economic growth render this WTU vulnerable. In North America, the  
 185 vulnerabilities are related to population growth and temperature increase.



186  
 187 **Figure 3: The vulnerability and projected change of the top 5 WTUs of each continent. The total vulnerability (indicated**  
 188 **by larger polygons), and projected change indicators of the top 5 important WTUs on each continent. BWS is the baseline**  
 189 **water stress indicator of the basin<sup>38</sup>; GE is an indicator for government effectiveness in the basin<sup>39</sup>; HT is hydro-political**  
 190 **tension<sup>37</sup>; dGDP<sup>41</sup> and dPop<sup>7</sup> are the projected changes in gross domestic product and population between 2000 and**  
 191 **2050, according to Shared Socioeconomic Pathway 2; dP<sup>40</sup> and dT<sup>40</sup> are the projected precipitation and temperature**  
 192 **changes between 2000 and 2050 according to the CMIP5 multi-model ensemble mean for Representative Concentration**  
 193 **Pathway 4.5. WTUs are ranked by vulnerability (highest vulnerability on top); colour filling indicates the WTU's Water**  
 194 **Tower Index (WTI) value. See Methods for calculation details.**

195



## **Water towers as global assets with increasing importance**

Planetary boundaries (PB) (e.g., the CO<sub>2</sub> concentration, global fresh water use, biosphere integrity) are defined as thresholds within which humanity can safely function without abrupt large scale changes to the environment<sup>43</sup>. Climate change and biosphere integrity have been identified as the core PBs with the potential to change the state of the Earth System should they be consistently transgressed for a prolonged period of time<sup>44</sup>. The global food system, in particular, has been identified as a major pressure on the PBs<sup>45</sup>. Without targeted technological changes and mitigation measures, it is expected that the adverse environmental effects of the food system could increase by more than 50% by 2050 relative to 2010, thus crossing the PBs<sup>45</sup>. In relation to the PBs, water towers are of particular importance. They are highly vulnerable to climate change, a key water supply that sustains the major global food systems in the world, and rich in biodiversity.

A clear implication is that vulnerability can be decreased with conservation, or increased with inefficient water use. This may seem logical and obvious, but it also means that the priorities for the most urgent action can be shifted as the nations of WTUs practice conservation or grow in an unsustainable way. Water conservation is the one part of the equation that is more or less under an individual nation's control. Efficient use of scarce water resources can translate into improved well-being of people and increased economic and food security.

The vulnerability of these water towers in the future is controlled by the trajectory of change that a WTU and its associated downstream basin will follow. At the global scale we made a first order assessment for a middle of the road scenario both in terms of climate change and socio-economic pathway (see Methods). However, it is important to acknowledge that the future pathways are highly precarious and the outcomes diverging and uncertain. A recent assessment for the Hindu-Kush Himalayan region concluded that there is no single likely future, and the region may run *downhill*, may do *business as usual* or it may advance to *prosperity*<sup>46</sup>. Each of those future pathways will result

in systematically different demands for water and may cross the PBs in varying degrees and this will likely hold for most WTUs in particular in Asia and South America.




Figure 4: Water Tower Index and vulnerabilities of the Indus basin. Panel A shows the supply and demand indicators and panel B the vulnerabilities. See Methods for details on the supply and demand indicators and the meaning of the vulnerability ranges.

227 Mountains are also an essential resource in the context of the UN Sustainable Development Goals  
228 (SDGs) that have been targeted towards the year 2030<sup>47</sup>. Mountains play a key role in achieving the  
229 SDGs for water (SDG 6), food (SDG 2) and energy (SDG 7). Given the projected change in climate and  
230 socioeconomic development in mountain-dependent basins, it is evident that if the SDGs are to be  
231 achieved the water resources of the Water Towers need to be harnessed within safe environmental  
232 limits.

233 It is essential to (i) recognize mountain regions as a global asset of the Earth system, (ii) acknowledge  
234 that vulnerability of the world's water towers is driven both by socio-economic factors and climate  
235 change and (iii) to develop international, mountain-specific conservation and climate change  
236 adaptation policies (e.g. national parks, pollutants control, emission reductions, erosion control, dam  
237 regulations) that safeguard the mountain ecosystems and mountain people and simultaneously  
238 ensure water, food and energy security of the millions of people downstream.

239

## Methods

### Delineation of Water Tower Units

In this study, we define a water tower unit (WTU) as the intersection of major river basins<sup>3</sup> and a topographic mountain classification based on elevation and surface roughness developed in the framework of the Global Mountain Biodiversity Assessment (GMBA)<sup>4</sup>. Although other similar mountain classification datasets exist<sup>1</sup>, that are also based on a combination of elevation and surface roughness, we use the GMBA classification (version 1.2) because topographical names of mountain ranges have been assigned to each of the mountain regions classified. The original GMBA inventory contains 1048 mountain regions worldwide. We make a subset of this dataset by imposing minimum thresholds for glacier area, glacier ice volume and snow persistence. We retain those mountain regions which have an ice volume larger than  $0.1 \text{ km}^3$ <sup>48</sup> or an average annual areal snow persistence larger than 10%<sup>5</sup>. After imposing these thresholds, 174 mountain regions remain. We intersect those regions with the major river basins and dissolve the result based on major river basin ID; i.e., all selected GMBA regions within a basin are grouped as a single WTU (Extended Data Figure 1, Extended Data Table 1, Extended Data Table 2). The final WTU delineation contains 78 units (Extended Data Figure 1). For each WTU we also define the downstream area that directly depends on the WTU using the river sub-basin delineation<sup>3</sup>, and we specify which mountain ranges are part of the WTU (Extended Data Figure 1, Extended Data Table 1, Extended Data Table 2). This dependent downstream area is smaller than the total downstream basin since not every downstream sub-basin is hydrologically connected to the WTU. To this end we start at the WTU and iteratively select each connected downstream sub-basin until the basin outlet, or lowest sub-basin in case of an endorheic system, is reached (Extended Data Figure 1).

### Quantifying the Water Tower Index

We combine a supply index and a demand index into a Water Tower Index (WTI) to rank WTUs. All grid calculations are performed at a  $0.05^\circ$  resolution.

The supply index (SI; Extended Data Table 3 for all equations) is based on indicators for precipitation, snow cover, glaciers, and surface water storage. For the precipitation indicator, the 2019 released ERA5 reanalysis dataset is used<sup>32</sup>. As sub-indicators, first, we compute the total annual average (2001-2017) WTU precipitation (Extended Data Figure 3a) relative to the overall basin precipitation ( $P_T$ ). We then include the inter-annual variation in WTU precipitation ( $P_{YV}$ ) and the intra-annual monthly WTU variation ( $P_{MV}$ ) based on the 2001-2017 time-series. We combine these three sub-indicators in a precipitation indicator ( $P$ ) where the variation ( $P_{YV}$  and  $P_{MV}$ ) has equal weight as  $P_T$ . The underlying assumption of including the variation is that if the variation is low, the WTU will provide a constant flow of water to the downstream basin, and therefore they are a more important WTU. For the snow cover indicator, we use the MODIS MOD10CM1 product<sup>5</sup>. We derive an average annual snow cover ( $S_T$ ) in each WTU for the 2001-2017 period (Extended Data Figure 3b). Here too, we derive both an inter-annual ( $S_{YV}$ ) and intra-annual variation ( $S_{MV}$ ) in snow cover, and using the same rationale as for the precipitation indicator, we combine the average snow persistence with the variation to derive a final snow indicator ( $S$ ). For the glacier indicator, we compute the glacier ice volume in a WTU<sup>48</sup> (Extended Data Figure 4a) relative to the average annual WTU precipitation ( $G_S$ ). We also compute the annual glacier melt water flux relative to the WTU precipitation on non-glacierised terrain ( $G_M$ ). We estimate the glacier melt water flux by the sum of the on-glacier precipitation and the mass balance per WTU. The WTU mass balance is based on the area weighted average annual mass balance from all geodetic and direct mass balance measurements made available by the World Glacier Monitoring Service<sup>49</sup>. However if there are less than 10 glaciers with data available within a WTU then the regional average is used<sup>17</sup>. We sum  $G_S$  and  $G_M$  to derive a final glacier indicator ( $G$ ). For the surface water indicator ( $L$ ), we compute the total volume of water that is stored in lakes and reservoirs in a WTU<sup>50</sup> (Extended Data Figure 4b) relative to the average annual WTU precipitation. The supply indicator (SI) is the average of  $P$ ,  $S$ ,  $G$  and  $L$ .

The demand index (DI) is based on net human water demands for domestic, industrial and irrigation purposes<sup>33</sup>, and natural demand (Extended Data Table 4 for all equations, Extended Data Figure 5,

Extended Data Figure 6). Since the natural demand, defined as the minimum river flow required to sustain the ecosystem, is not readily available, we estimate this with the environmental flow requirement computed with the 90<sup>th</sup> percentile exceedance value of the natural flow<sup>33,51,52</sup>. First, the average monthly sectoral demands are computed based on a 2001-2014 time series ( $D_{DOM,m}$ ,  $D_{IRR,m}$ ,  $D_{IND,m}$ ,  $D_{NAT,m}$ ). Part of each sectoral demand can potentially be met by downstream water availability that does not have its origin in the mountains. For each grid cell with a positive demand we therefore compute the average monthly water availability ( $WA_{DOM,m}$ ,  $WA_{IRR,m}$ ,  $WA_{IND,m}$ ,  $WA_{NAT,m}$ ; see Extended Data Table 4) as the precipitation minus the actual natural evapotranspiration (P-ET)<sup>32</sup>. We subtract this amount from the average monthly sectoral water demands as an estimate for the monthly demand that needs to be met by other sources, including the WTUs. We assume that the entire water deficit has to be provided by the WTU, although other water sources, such as groundwater<sup>51</sup>, can also be important. We acknowledge that the global scale of our assessment also limits taking into account the distribution and allocation of water within different portions of our spatial units of calculation. Finally, we aggregate these monthly net demands to be sustained by the WTU over all months and we divide it by the total annual sectoral demand to get four demand indicators ( $D_{DOM}$ ,  $D_{IND}$ ,  $D_{IRR}$ ,  $D_{NAT}$ ). The demand index (DI) is the average of the indicators  $D_{DOM}$ ,  $D_{IND}$ ,  $D_{IRR}$ , and  $D_{NAT}$ .

The final Water Tower Index (WTI) is the product of SI and DI, for which the values are subsequently normalized over the range of WTI values found for all 78 WTUs. By using a multiplicative approach, we ensure that a WTU only ranks high when it has considerable water resources (either as precipitation, glacier ice, snow and surface water or a combination) in the mountains, and the demand for those resources downstream is likewise high (Extended Data Figure 2).

## Uncertainty

It is acknowledged the SI, DI and WTI are based on partly arbitrary choices of indicators and sub-indicators. In our assessment we have assigned an equal weight to each of the indicators constituting SI and DI. To account for uncertainty in the weight of each indicator in the WTI calculation we have

performed a sensitivity analysis where we randomly vary the weights of each of the eight indicators that constitute the SI and DI and assess the impact on the WTI-ranking of the WTUs. We assume that the weight of each indicator is uniformly distributed and can be a maximum of three times as high or low as another indicator, and we assess through a 10,000 member Monte Carlo analysis how sensitive the rank of the WTU is as a result of this uncertainty. The analysis shows that the top and bottom of the ranking are robust and only limited shifts in the ranking occur (< 5 positions). However, the middle part of the ranking is more sensitive to the weights of the indicators and there is a considerable number of WTUs where in more than 25% of the total runs the rank changes more than 5 positions.

In addition, we also include a 1000 member Monte Carlo analysis to assess the propagation of uncertainty in the datasets used in the WTI calculation. For each input dataset we estimate a standard deviation and assuming a normally distributed error we sample from the distribution to assess how the input data uncertainty affects the WTI value (Table S1) and WTU ranking (Extended Data Figure 7). For precipitation we compute the standard deviation per WTU and per downstream basin based on 9 different precipitation datasets (CRU bias-corrected with ERA-Interim, CRU TS2.1 downscaled with ERA-40, CRU TS3.21 downscaled with ERA-40, CRU TS3.21 downscaled with ERA-Interim, WFDEI, NCEP-NCAR Reanalysis, WATCH, WATCH corrected with GPCC, ERA5)<sup>32,53–59</sup>. For evapotranspiration we take a similar approach using 4 different datasets (ERA-Interim, GLEAM, MERRA-2, PCR-GLOBWB forced with ERA-Interim, ERA5)<sup>32,54,60–62</sup>. Values for snow persistence, ice volumes, glacier mass balance, and the domestic, industrial and irrigation water demands are derived from literature<sup>17,48,63–65</sup>. For the uncertainty in lake and reservoir volume we assume a standard deviation of 10% and we keep the environmental flow requirement constant. The ranking is also sensitive to input data uncertainty; however, the ranking is robust in particular in the top 20 of the ranking where only limited shifts in positions occur. Here, too, most shifts are observed in the middle part of the ranking.

The code and data used in this uncertainty analysis is provided online (see Data and Code Availability Section).

### **Assessing vulnerabilities**

For the WTUs, we assess the vulnerability of their role as water tower based on three static indicators for water stress, government effectiveness, and the potential for hydro-political tension in case of transboundary basins (Table S2). In addition, we include four change indicators: the projected change in temperature, precipitation, population and gross domestic product between 2000 and 2050. In all cases we use the ensemble mean RCP4.5 climate change scenario<sup>66</sup> in combination with SSP2 shared socio-economic pathway<sup>67</sup> as a middle of the road scenario, both in terms of economic development and associated climate change (Table S2). We scale the different vulnerability indicators between 0 (minimum vulnerability) and 1 (maximum vulnerability) considering the thresholds defined below.

For water stress, we use the Baseline Water Stress (BWS) indicator<sup>38</sup>. BWS measures the ratio of total water withdrawals to available renewable surface and groundwater supplies with higher values indicating more competition among users. The index value is derived from an ordinary least squares regression fitted through raw monthly water stress values for 1960-2014, taking the fitted BWS value for 2014<sup>38</sup>. We compute the area-averaged BWS for all WTUs including their downstream dependent areas and scale between 0 and 5 which is the range of the BWS scale in the cited study. High BWS is associated to high vulnerability and low BWS is associated with low vulnerability. Since no global dataset for water management capacity is available at global scale we validated the indicators Gross Domestic Product (GDP)<sup>68</sup>, Human Development Index (HDI)<sup>68</sup> and Government Effectiveness (GE)<sup>39</sup> as proxies for water management capacity, which is available for selected mountainous basins only<sup>14</sup>. GE shows best correlation to water management capacity in the selection of basins, and we calculate the area-averaged value for each WTU including its downstream dependent area. We scale between -1.5 and 2.0 which are the minimum and maximum values found for the WTUs. A low value for GE



implies high vulnerability while a high value for GE indicates low vulnerability. Lastly, all transboundary basins are assessed on the risk for potential hydro-political tensions that are based on a global mapping of basins that are ill-equipped to deal with transboundary disputes triggered by the construction of new dams and diversions<sup>37</sup>. We compute the WTU basin aggregated score provided by the cited study and the range of the original scale in the cited study (0-5) is used to scale between minimum and maximum.

For each WTU we compute a projected multi-model ensemble mean change in precipitation (%) and temperature (K) between 2000 and 2050 for RCP4.5 for 35 different CMIP5 climate models<sup>40</sup>. For projected changes in temperature the scores for the individual WTUs are linearly scaled between 0 and 1 for the full range of projected temperature increases of all WTUs. For precipitation projections, only decreases in precipitation are assumed to contribute to vulnerability (i.e. projections of increases in precipitation and unchanged precipitation are classified as minimum vulnerability). The scores for the individual WTUs are scaled linearly between 0 and 1, where 0 indicates unchanged or increasing precipitation and 1 indicates the largest precipitation decrease projected for all 78 WTUs. The projected population change between 2016 and 2050 for Shared Socioeconomic Pathway (SSP) 2 is derived from the HYDE database<sup>7</sup> and the relative increase for each of the WTU basins is computed. All WTUs are scaled between a growth of 0% and a maximum of 50%, i.e. if the projected population growth is more than 50%, a WTU has maximum vulnerability. The relative increase in GDP between 2000 and 2050 is computed per WTU basin, with the assumption that a strong projected increase in GDP is indicative of a strong growth in water demand. Data for the SSP2 shared socio-economic pathway are used<sup>41</sup>. All WTU basins are scaled between the minimum and the maximum, which is capped by a growth rate of 1000%.

We assess indicators of various nature for vulnerability and future changes. To assess a complete vulnerability based on this set of indicators is challenging and requires knowledge of the weights of the individual indicators in assessing the total vulnerability for each WTU. The caveat is made that we

consider a middle of the road scenario both in terms of projected climate change and socio-economic development as a first order assessment. The future development pathway in most WTUs, in particular in Asia and South America, is uncertain and highly diverging and depends on the global economy, regional growth rates and geopolitical tensions, which are difficult to project or quantify. In addition, satisfactory representation of mountainous climate in General Circulation Models is difficult, leading to large uncertainty in particular for future precipitation projections.

In our study we assess impacts-driven vulnerability, where vulnerability is defined in direct proportion to the magnitude of hydrological change. However, we note that recent work on the human dimensions of climate change have demonstrated that vulnerability emerges from the interaction of both environmental and social dynamics in specific contexts<sup>69,70</sup>.

## **Acknowledgements**

This project was funded by a partnership between National Geographic Society and Rolex, through the Life at the Extremes Initiative. The authors are grateful to the Strategic Priority Research Program of the Chinese Academy of Sciences for their support, to Daniel Farinotti for providing the data on glacier volume, and to Niko Wanders for providing precipitation datasets used in the uncertainty analysis

## **Author Contributions**

W.W.I. and A.F.L. contributed equally to the study; they designed the study, performed the analysis, prepared figures and tables and drafted the manuscript. P.D.A.K. contributed to the data analysis and prepared Figure 3. Y.W. provided the dataset used to calculate demand indicators. S.B., S.H., A.B. and A.C.E contributed to the design of the index and analysis methods. All authors contributed to developing the theory and conception of the study by providing regional (M.A., A.F., and P.P. for the Andes; T.B. U.H., P.D.A.K., A.V.K., P.M., S.N, F.P., A.B.S., A.S., C.X., and T.Y. for High Mountain Asia; T.B., A.E., F.P., and D.V. for the Alps; and S.R., T.P., J.S.K., M.K. for North America) and/or subject specific expertise (B.J.D., J.S.K., A.B.S., P.P., A.S and S.R. for glacial volume; U.H., M.K. and F.P. for

meltwater discharge; H.B., A.F. and Y.W. on irrigation demand; T.B., A.E., J.S.K. and A.V.K. for glacial lakes; M.F. and T.H.P. for global snow cover, P.D.A.K. for volume ice loss; A.F., P.M., A.S. and T.Y. for climatology; S.N. and S.R. for hydrology; M.K., A.B.S. and D.V. for water demand, conflicts, and vulnerability; H.R. for preferential flow; S.R. for glacier accumulation mass loss and its effects on downstream populations; D.V. for water management capacity; C.X. for global cryospheric functions and processes; and Y.W. for environmental flow requirements). All authors discussed and provided feedback on the manuscript. The study was initiated by J.E.M.B. and facilitated by A.C.E.

## **Data availability**

### *Data generated in this study*

The data generated to support the findings of this study are available in an online data repository at zenodo.org with a doi identifier upon acceptance of the manuscript. The data is temporarily available during peer-review at

[https://www.dropbox.com/s/3sbrkmqp960btik/data\\_ngs\\_watertowers.zip?dl=0](https://www.dropbox.com/s/3sbrkmqp960btik/data_ngs_watertowers.zip?dl=0).

### *Third party data used in this study*

- Hydrological basin boundaries<sup>3</sup> used in this study are available online at <http://www.fao.org/nr/water/aquamaps/>.
- Mountain definition data<sup>4</sup> used in this study are available online at [https://ilias.unibe.ch/goto\\_ilias3\\_unibe\\_file\\_1047348.html](https://ilias.unibe.ch/goto_ilias3_unibe_file_1047348.html).
- Precipitation data used in this study<sup>32</sup> are available online at <https://cds.climate.copernicus.eu>.
- Snow cover data used in this study<sup>5</sup> are available online at <https://nsidc.org/data/mod10cm>.
- Glacier volume data<sup>48</sup> used in this study are available online at <https://doi.org/10.3929/ethz-b-000315707>.
- Glacier mass balance data<sup>17,71</sup> are available online at <https://wgms.ch/>.

- Lake and reservoir storage data<sup>50</sup> used in this study are available online at  
<https://www.hydrosheds.org/pages/hydrolakes>.
- Water demand data used in this study are available upon request from Y. Wada  
(wada@iiasa.ac.at).
- Baseline Water Stress data<sup>72</sup> used in this study are available online at  
<https://www.wri.org/aqueduct>.
- Government Effectiveness data<sup>39</sup> used in this study are available online at  
<https://info.worldbank.org/governance/wgi/#home>.
- Data on hydro-political tensions for transboundary river basins<sup>37</sup> used in this study are  
available online at  
<https://transboundarywaters.science.oregonstate.edu/content/transboundary-freshwater-spatial-database>.
- Data for future projections of population count<sup>7</sup> used in this study are available online at  
<ftp://ftp.pbl.nl/hyde/SSPs/SSP2/zip/>.
- Data for future projections of Gross Domestic Product<sup>41</sup> used in this study are available  
online at <http://www.cger.nies.go.jp/gcp/population-and-gdp.html>.
- Data for future projections of temperature and precipitation<sup>40</sup> used in this study are available  
online at <https://climexp.knmi.nl>.

## Code Availability

All code developed for the calculations performed for this study are publicly available in a Github repository at [https://github.com/mountainhydrology/pub\\_ngo-watertowers](https://github.com/mountainhydrology/pub_ngo-watertowers) upon acceptance of the manuscript. A copy of the repository is temporarily available during peer-review at [https://www.dropbox.com/s/ktg55kg6b1mug7a/pub\\_ngo-watertowers-master.zip?dl=0](https://www.dropbox.com/s/ktg55kg6b1mug7a/pub_ngo-watertowers-master.zip?dl=0)

## References

1. Viviroli, D., Dürr, H. H., Messerli, B., Meybeck, M. & Weingartner, R. Mountains of the world, water towers for humanity: Typology, mapping, and global significance. *Water Resour. Res.* **43**, 1–13 (2007).
2. Immerzeel, W. W., Van Beek, L. P. H. & Bierkens, M. F. P. Climate change will affect the Asian water towers. *Science* **328**, 1382–1385 (2010).
3. Lehner, B., Verdin, K. & Jarvis, A. New global hydrography derived from spaceborne elevation data. *Eos Trans. AGU* **89**, 93–94 (2008).
4. Körner, C. *et al.* A global inventory of mountains for bio-geographical applications. *Alp. Bot.* **127**, 1–15 (2017).
5. Hall, D. K. & Riggs, G. A. MODIS/Terra Snow Cover Monthly L3 Global 0.05Deg CMG, Version 6. (2015). doi:10.5067/MODIS/MOD10CM.006
6. Pfeffer, W. T. *et al.* The Randolph Glacier Inventory: a globally complete inventory of glaciers. *J. Glaciol.* **60**, 537–552 (2014).
7. Goldewijk, K. K., Beusen, A. & Drecht, G. Van. The HYDE 3 . 1 spatially explicit database of human-induced global land-use change over the past 12,000 years. *Glob. Ecol. Biogeogr.* **20**, 73–86 (2011).
8. Xiao, C. De, Wang, S. J. & Qin, D. H. A preliminary study of cryosphere service function and value evaluation. *Adv. Clim. Chang. Res.* **6**, 181–187 (2015).
9. Wang, X., Liu, S. W. & Zhang, J. L. A new look at roles of the cryosphere in sustainable development. *Adv. Clim. Chang. Res.* **10**, 124–131 (2019).
10. Chape, S., Spalding, M. D. & Jenkins, M. D. *The world's protected areas.* (UNEP-World

- 487 Conservation Monitoring Centre, 2008).
- 488 11. Körner, C. & Paulsen, J. A world-wide study of high altitude treeline temperatures. *J. Biogeogr.*  
489 **31**, 713–732 (2004).
- 490 12. Körner, C., Paulsen, J. & Spehn, E. M. A definition of mountains and their bioclimatic belts for  
491 global comparisons of biodiversity data. *Alp. Bot.* **121**, 73–78 (2011).
- 492 13. Nordhaus, W. D. Geography and macroeconomics : New data and new findings. (2006).
- 493 14. Viviroli, D. *et al.* Climate change and mountain water resources: overview and  
494 recommendations for research, management and policy. *Hydrol. Earth Syst. Sci.* **15**, 471–504  
495 (2011).
- 496 15. IPCC. *Special Report: The Ocean and Cryosphere in a Changing Climate.*  
497 doi:<https://www.ipcc.ch/report/srocc/>
- 498 16. Pepin, N. *et al.* Elevation-dependent warming in mountain regions of the world. *Nat. Clim.*  
499 *Chang.* **5**, 424–430 (2015).
- 500 17. Zemp, M. *et al.* Global glacier mass changes and their contributions to sea-level rise from  
501 1961 to 2016. *Nature* **568**, 382–386 (2019).
- 502 18. Hammond, J. C., Saavedra, F. A. & Kampf, S. K. Global snow zone maps and trends in snow  
503 persistence 2001-2016. *Int. J. Climatol.* 1–15 (2018). doi:10.1002/joc.5674
- 504 19. Bormann, K. J., Brown, R. D., Derksen, C. & Painter, T. H. Estimating snow-cover trends from  
505 space. *Nat. Clim. Chang.* (2018). doi:10.1038/s41558-018-0318-3
- 506 20. Sarangi, C. *et al.* Impact of light-absorbing particles on snow albedo darkening and associated  
507 radiative forcing over High Mountain Asia: High resolution WRF-Chem modeling and new  
508 satellite observations. *Atmos. Chem. Phys. Discuss.* 1–56 (2018). doi:10.5194/acp-2018-979

- 509 21. Painter, T. H. *et al.* Impact of disturbed desert soils on duration of mountain snow cover.  
510 *Geophys. Res. Lett.* **34**, 1–6 (2007).
- 511 22. Lutz, A. F., Immerzeel, W. W., Shrestha, A. B. & Bierkens, M. F. P. Consistent increase in High  
512 Asia ' s runoff due to increasing glacier melt and precipitation. *Nat. Clim. Chang.* **4**, 1–6 (2014).
- 513 23. Kargel, J. S. S. *et al.* Geomorphic and geologic controls of geohazards induced by Nepals 2015  
514 Gorkha earthquake. *Science (80-. )*. **351**, 140 (2016).
- 515 24. Kirschbaum, D. *et al.* The State of Remote Sensing Capabilities of Cascading Hazards Over High  
516 Mountain Asia. *Front. Earth Sci.* **7**, (2019).
- 517 25. Guha-Sapir, D., Below, R. & Hoyois, P. *EM-DAT: International Disaster Database*. (2019).
- 518 26. Mal, S. *Climate Change , Extreme Events and Disaster Risk Reduction*. (Springer, 2018).
- 519 27. Mann, M. E. *et al.* Influence of Anthropogenic Climate Change on Planetary Wave Resonance  
520 and Extreme Weather Events. *Sci. Rep.* **7**, (2017).
- 521 28. Fischer, E. M. & Knutti, R. Anthropogenic contribution to global occurrence of heavy-  
522 precipitation and high-temperature extremes. *Nat. Clim. Chang.* **5**, 560–564 (2015).
- 523 29. Haeberli, W., Schaub, Y. & Huggel, C. Increasing risks related to landslides from degrading  
524 permafrost into new lakes in de-glaciating mountain ranges. *Geomorphology* **293**, 405–417  
525 (2017).
- 526 30. Huss, M. *et al.* Toward mountains without permanent snow and ice. *Earth's Futur.* **5**, 418–435  
527 (2017).
- 528 31. Biemans, H. *et al.* Importance of snow and glacier meltwater for agriculture on the Indo-  
529 Gangetic Plain. *Nat. Sustain.* **2**, 594–601 (2019).
- 530 32. C3S. ERA5: Fifth generation of ECMWF atmospheric reanalyses of the global climate. (2017).

- 531 33. Wada, Y., De Graaf, I. E. M. & van Beek, L. P. H. High-resolution modeling of human and  
532 climate impacts on global water resources. *J. Adv. Model. Earth Syst.* **8**, 735–763 (2016).
- 533 34. Wada, Y. *et al.* Global depletion of groundwater resources. *Geophys. Res. Lett.* **37**, 1–5 (2010).
- 534 35. Wada, Y., van Beek, L. P. H. & Bierkens, M. F. P. Nonsustainable groundwater sustaining  
535 irrigation: A global assessment. *Water Resour. Res.* **48**, (2012).
- 536 36. Immerzeel, W. W. & Bierkens, M. F. P. Asia’s water balance. *Nat. Geosci.* **5**, 841–842 (2012).
- 537 37. De Stefano, L., Petersen-Perlman, J. D., Sproles, E. A., Eynard, J. & Wolf, A. T. Assessment of  
538 transboundary river basins for potential hydro-political tensions. *Glob. Environ. Chang.* **45**,  
539 35–46 (2017).
- 540 38. Hofste, R. W. *et al.* *Aqueduct 3.0: Updated decision-relevant global water risk indicators.*  
541 *Technical Note.* (2019).
- 542 39. Kaufmann, D., Kraay, A. & Mastruzzi, M. *The Worldwide Governance Indicators. Methodology*  
543 *and Analytical Issues. Policy Research Working Paper 5430* (2010).
- 544 40. Taylor, K. E., Stouffer, R. J. & Meehl, G. A. An overview of CMIP5 and the experiment design.  
545 *Bull. Am. Meteorol. Soc.* **93**, 485–498 (2012).
- 546 41. Murakami, D. & Yamagata, Y. Estimation of gridded population and GDP scenarios with  
547 spatially explicit statistical downscaling. *ArXiv* **1610.09041**, (2016).
- 548 42. Wijngaard, R. R., Biemans, H., Lutz, A. F., Shrestha, A. B. & Immerzeel, W. W. Climate change  
549 vs. Socio-economic development: Understanding the South-Asian water gap. *Hydrol. Earth*  
550 *Syst. Sci* **22**, 6297–6321 (2018).
- 551 43. Rockström, J. *et al.* A safe operating space for humanity. *Ecol. Soc.* **14**(2) **14**, (2009).
- 552 44. Steffen, W. *et al.* Planetary boundaries: Guiding human development on a changing planet.



- 553        *Science* (80-. ). **347**, 1259855 (2015).
- 554    45.    Springmann, M. *et al.* Options for keeping the food system within environmental limits.  
555        *Nature* **562**, 519–525 (2018).
- 556    46.    Roy, J. *et al.* Exploring Futures of the Hindu Kush Himalaya: Scenarios and Pathways. in *The*  
557        *Hindu Kush Himalaya Assessment: Mountains, Climate Change, Sustainability and People* (eds.  
558        Wester, P., Mishra, A., Mukherji, A. & Shrestha, A. B.) 99–125 (Springer International  
559        Publishing, 2019). doi:10.1007/978-3-319-92288-1\_4
- 560    47.    UN. Transforming our world: the 2030 Agenda for Sustainable Development, A/RES/70/1.  
561        (2015).
- 562    48.    Farinotti, D. *et al.* A consensus estimate for the ice thickness distribution of all glaciers on  
563        Earth. *Nat. Geosci.* **12**, 168–173 (2019).
- 564    49.    World Glacier Monitoring Service. Fluctuations of Glaciers (FoG) Database. (2018).  
565        doi:10.5904/wgms-fog-2018-06
- 566    50.    Messenger, M. L., Lehner, B., Grill, G., Nedeva, I. & Schmitt, O. Estimating the volume and age  
567        of water stored in global lakes using a geo-statistical approach. *Nat. Commun.* **7**, 1–11 (2016).
- 568    51.    Gleeson, T., Wada, Y., Bierkens, M. F. P. & van Beek, L. P. H. Water balance of global aquifers  
569        revealed by groundwater footprint. *Nature* **488**, 197–200 (2012).
- 570    52.    Smakhtin, V., Revenga, C. & Döll, P. A pilot global assessment of environmental water  
571        requirements and scarcity. *Water Int.* **29**, 307–317 (2004).
- 572    53.    Harris, I., Jones, P. D., Osborn, T. J. & Lister, D. H. Updated high-resolution grids of monthly  
573        climatic observations - the CRU TS3.10 Dataset. *Int. J. Climatol.* (2013). doi:10.1002/joc.3711
- 574    54.    Dee, D. P. *et al.* The ERA-Interim reanalysis: configuration and performance of the data

575 assimilation system. *Q. J. R. Meteorol. Soc.* **137**, 553–597 (2011).

576 55. Weedon, G. P. *et al.* The WFDEI meteorological forcing data set: WATCH Forcing Data  
577 methodology applied to ERA-Interim reanalysis data. *Water Resour. Res.* 7505–7514 (2014).  
578 doi:10.1002/2014WR015638. Received

579 56. Weedon, G. P. *et al.* Creation of the WATCH Forcing Data and Its Use to Assess Global and  
580 Regional Reference Crop Evaporation over Land during the Twentieth Century. *J.*  
581 *Hydrometeorol.* **12**, 823–848 (2011).

582 57. Schneider, U. *et al.* GPCC's new land surface precipitation climatology based on quality-  
583 controlled in situ data and its role in quantifying the global water cycle. *Theor. Appl. Climatol.*  
584 26 (2013). doi:10.1007/s00704-013-0860-x

585 58. Uppala, S. M. *et al.* The ERA-40 re-analysis. *Q. J. R. Meteorol. Soc.* **131**, 2961–3012 (2005).

586 59. Kalnay, E. *et al.* The NCEP/NCAR 40-Year Reanalysis Project. *Bull. Am. Meteorol. Soc.* **77**, 437–  
587 471 (1996).

588 60. Martens, B. *et al.* GLEAM v3: Satellite-based land evaporation and root-zone soil moisture.  
589 *Geosci. Model Dev.* **10**, 1903–1925 (2017).

590 61. Rienecker, M. M. *et al.* MERRA: NASA's Modern-Era Retrospective Analysis for Research and  
591 Applications. *J. Clim.* **24**, 3624–3648 (2011).

592 62. Sutanudjaja, E. H. *et al.* PCR-GLOBWB 2: A 5 arcmin global hydrological and water resources  
593 model. *Geosci. Model Dev.* **11**, 2429–2453 (2018).

594 63. Riggs, G. A., Hall, D. K. & Román, M. O. Overview of NASA's MODIS and VIIRS Snow-Cover  
595 Earth System Data Records. *Earth Syst. Sci. Data* 1–30 (2017). doi:10.5194/essd-2017-25

596 64. Wada, Y. *et al.* Modeling global water use for the 21st century: The Water Futures and

- 597 Solutions (WFaS) initiative and its approaches. *Geosci. Model Dev.* **9**, 175–222 (2016).
- 598 65. Wada, Y. *et al.* Multimodel projections and uncertainties of irrigation water demand under  
599 climate change. *Geophys. Res. Lett.* **40**, 4626–4632 (2013).
- 600 66. van Vuuren, D. P. *et al.* The representative concentration pathways: an overview. *Clim.*  
601 *Change* **109**, 5–31 (2011).
- 602 67. O’Neill, B. C. *et al.* The roads ahead: Narratives for shared socioeconomic pathways describing  
603 world futures in the 21st century. *Glob. Environ. Chang.* **42**, 169–180 (2017).
- 604 68. Kummu, M., Taka, M. & Guillaume, J. H. A. Gridded global datasets for Gross Domestic  
605 Product and Human Development Index over 1990–2015. *Sci. Data* **5**, 1–15 (2018).
- 606 69. McDowell, G. *et al.* Adaptation action and research in glaciated mountain systems: Are they  
607 enough to meet the challenge of climate change? *Glob. Environ. Chang.* **54**, 19–30 (2019).
- 608 70. Conway, D. *et al.* The need for bottom-up assessments of climate risks and adaptation in  
609 climate-sensitive regions. *Nat. Clim. Chang.* **9**, (2019).
- 610 71. Zemp, M. *et al.* *WGMS (2017): Global Glacier Change Bulletin No. 2 (2014–2015)*. (2017).
- 611 72. Gassert, F., Landis, M., Luck, M., Reig, P. & Shiao, T. *Aqueduct Global Maps 2.1. World*  
612 *Resources Institute Working Paper* (2014).
- 613 73. Wada, Y., de Graaf, I. E. M. & van Beek, L. P. H. High-resolution modeling of human and  
614 climate impacts on global water resources. *J. Adv. Model. Earth Syst.* **8**, 735–763 (2016).

615

# Importance and vulnerability of the world's water towers

*Immerzeel, W.W., Lutz, A.F., Andrade, M., Bahl, A., Biemans, H., Bolch, T., Hyde, S., Brumby, S.,  
Davies, B.J., Elmore, A.C., Emmer, A., Feng, M., Fernández, A., Haritashya, U., Kargel, J.S., Koppes, M.,  
Kraaijenbrink, P.D.A., Kulkarni, A.V., Mayewski, P., Nepal, S., Pacheco, P., Painter, T.H., Pellicciotti, F.,  
Rajaram, H., Rupper, S., Sinisalo, A., Shrestha, A.B., Viviroli, D., Wada, Y., Xiao, C., Yao, T., Baillie,  
J.E.M.*

## Extended data figures and tables

## 628 Extended Data Tables

629 **Extended Data Table 1: List of WTUs and the GMBA mountain ranges which are (partly) covered by each WTU, for North**  
 630 **America and South America**

WTU ID	WTU Name	GMBA mountain ranges (partly) covered by WTU
1	Mississippi-Missouri	Bighorn Mountains, Absaroka Range, Crazy Mountains, Lewis Range, Swan Range, Flathead Range, Wind River Range, Front Range, Medicine Bow Mountains, Gore Range, Sawatch Range
2	California	Sierra Nevada, Cascade Range
3	Great Basin	Sierra Nevada
4	N-America, Colorado	San Juan Mountains, Wind River Range, Front Range, Medicine Bow Mountains, Gore Range, Sawatch Range
5	Columbia and NW-US	Absaroka Range, Lewis Range, Swan Range, Mission Range, Flathead Range, Purcell Mountains, Cabinet Mountains, Sawtooth Mountains, Teton Range, Wind River Range, Wallowa Mountains, Cariboo Mountains, Monashee Mountains, Selkirk Mountains, Coast Mountains, Rocky Mountains Calgary, Scrip Range, Cascade Range
6	Fraser	Coast Mountains, Skeena Mountains, Omineca Mountains, Cariboo Mountains, Monashee Mountains, Hazelton Mountains, Rocky Mountains Calgary, Scrip Range, Cascade Range
7	Pac. and Arc. Coast	Chugach Mountains, Kenai Mountains, Alaska Range, Coast Mountains, Aleutian Range, Kodiak and Afognak Island, Alexander Archipelago, Vancouver Island, Brooks Range, Saint Elias Mountains, Wrangell Mountains, Kilbuck Mountains, Talkeetna Mountains, Mackenzie Mountains, Wernecke Mountains, Selwyn Mountains, Pelly Mountains, Skeena Mountains, Stikine Ranges, Cassiar Mountains, Omineca Mountains, Hazelton Mountains, Cascade Range, Olympic Mountains
8	Saskatchewan-Nelson	Lewis Range, Rocky Mountains Calgary
9	NW-Territ. and Nunavut	Baffin Island
10	Hudson Bay Coast	Tornat Mountains
11	Atl. Ocean Seaboard	Tornat Mountains
12	Mackenzie	Mackenzie Mountains, Wernecke Mountains, Selwyn Mountains, Pelly Mountains, Stikine Ranges, Cassiar Mountains, Omineca Mountains, Rocky Mountains Calgary
13	Río Grande-Bravo	San Juan Mountains, Sawatch Range
14	Caribbean Coast	Sierra Nevada de Santa Marta, Cordillera Oriental Colombia Venezuela
15	Magdalena	Sierra Nevada de Santa Marta, Cordillera Central Colombia, Cordillera Oriental Colombia Venezuela
16	Orinoco	Cordillera Oriental Colombia Venezuela
17	Amazon	Cordillera Central Colombia, Cordillera Oriental Colombia Venezuela, Cordillera Central Ecuador, Cordillera Oriental Peru Bolivia, Cordillera Occidental Peru Bolivia Chile, Altiplano
18	La Plata	Cordillera Oriental Peru Bolivia
19	S-America, Colorado	Cordillera principal, Cordillera de Oliva, Cordillera de Ollita, Cerro de Ansilta, Central Volcanic Zone, Cordillera Frontal
20	Negro	Cordillera principal, Northern Patagonian Andes
21	S-Arg., S-Atl. Coast	Cordillera Patagonica Sur, Northern Patagonian Andes, Andes fueginos
22	Col.-Ecuad., Pac. Coast	Cordillera Central Colombia, Cordillera Central Ecuador
23	Peru, Pac. Coast	Cordillera Central Ecuador, Cordillera Occidental Peru Bolivia Chile
24	N-Chile, Pac. Coast	Cordillera principal, Cordillera Occidental Peru Bolivia Chile, Sierra de la Punilla, Sierra de Tatul, Cordillera de Oliva, Cordillera de Ollita, Cerro de Ansilta, Central Volcanic Zone, Cordillera Frontal
25	S-Chile, Pac. Coast	Northern Patagonian Andes, Cordillera Patagonica Sur, Andes fueginos, Cordillera principal, Cordillera Frontal
26	La Puna Region	Cordillera Oriental Peru Bolivia, Cordillera Occidental Peru Bolivia Chile, Altiplano, Central Volcanic Zone, Cordillera Frontal
27	Salinas Grandes	Central Volcanic Zone, Cordillera Frontal

**Extended Data Table 2: List of WTUs and the GMBA mountain ranges which are (partly) covered by each WTU, for Europe, Asia and**

<b>Oceania</b>		
<b>WTU ID</b>	<b>WTU Name</b>	<b>GMBA mountain ranges (partly) covered by WTU</b>
28	Spain-Portugal, Atl. Coast	Pyrenees
29	Spain, S-and E-Coast	Pyrenees
30	Ebro	Pyrenees
31	Gironde	Pyrenees
32	France, W-Coast	Pyrenees
33	Rhône	European Alps
34	France, S-Coast	European Alps, Pyrenees
35	Rhine	European Alps
36	Po	European Alps, Pyrenees
37	Italy, W-Coast	European Alps
38	Italy, E-Coast	European Alps
39	Danube	European Alps, Dinaric Alps
40	Sweden	Scandinavian Mountains, Jotunheimen
41	Adr. Sea-Bl. Sea Coast	Dinaric Alps, European Alps
42	Volga	Ural Mountains
43	Ural	Ural Mountains
44	Black Sea, N-Coast	Greater Caucasus
45	Caspian Sea Coast	Greater Caucasus
46	Scandinavia, N-Coast	Scandinavian Mountains, Jotunheimen
47	Russia, Barents Sea Coast	Ural Mountains
48	Arc. Ocean Islands	Svalbard, Greenland Kalaallit Nunaat, Novaya Zemlya
49	Iceland	Iceland
50	Gobi Int.	Haanhöhiy Uul, Borohoro-Shan, Khrebet Dzhungarskiy Alatau, Khrebet Saur, Bogda Shan, Karlik Shan, Tulai Nanshan, Tulai Shan, Zoulang Nanshan, Lenglong Ling, Datong Shan, Türgen Uul, Kuroyskiy Khrebet, Shopshal'skiy Khrebet, Altai Mountains, Tien Shan
51	Yellow River	Lenglong Ling, Datong Shan, Banyan Har Shan, Qionglai Shan, Anyemaqen Shan
52	Tarim Int.	Alayskiy Khrebet, Ferganskiy Khrebet, Terskey Ala Too, Kokshaal Too, Borohoro-Shan, Narat Shan, Horo Shan, Eren Habirga Shan, Danghe Nanshan, Qaidam Shan, Tergun Daba Shan, Yema Shan, Shule Nanshan, Tulai Nanshan, Datong Shan, Pamir, Karakorum, Banyan Har Shan, Anyemaqen Shan, Tibetan Plateau, Kunlun Shan, Tien Shan
53	Tibetan Plateau	Nganglong Kangri, Gangdise Shan, Nyainqentanglha Shan, Tanggula Shan, Tibetan Plateau
54	Yangtze	Tanggula Shan, Banyan Har Shan, Ningjing Shan, Chola Shan, Shaluli Shan, Daxue Shan, Qionglai Shan, Tibetan Plateau, Yun Range
55	Mekong	Tanggula Shan, Ningjing Shan, Patkai Hills, Mishmi Hills, Tibetan Plateau, Yun Range
56	Salween	Nyainqentanglha Shan, Tanggula Shan, Patkai Hills, Mishmi Hills, Tibetan Plateau
57	Irrawaddy	Patkai Hills
58	Ganges-Bramaputra	Gangdise Shan, Nyainqentanglha Shan, Tanggula Shan, Himalaya, Patkai Hills, Mishmi Hills
59	Sabarmati	Himalaya
60	Indus	Himalaya, Ladakh Range, Pamir Mountains, Karakorum, Hindu Kush, Nganglong Kangri, Gangdise Shan, Malakand Range, Tibetan Plateau
61	Lena	Baykal'skiy Khrebet, Khrebet Kodar, Verkhoyanskiy Khrebet, Khrebet Suntar Khayata
62	Siberia, N-Coast	Gory Putorana
63	Yenisey	Haanhöhiy Uul, Shopshal'skiy Khrebet, Kuznetskiy Alatau, Zapadnyy Sayan, Vostochnyy Sayan, Baykal'skiy Khrebet, Gory Putorana
64	Kara Sea Coast	Ural Mountains
65	Ob	Khrebet Saur, Seminskiy Khrebet, Aygulakskiy Khrebet, Kuroyskiy Khrebet, Shopshal'skiy Khrebet, Kuznetskiy Alatau, Zapadnyy Sayan, Altai Mountains, Ural Mountains
66	Siberia, W-Coast	Chukotskiy (Anadyrskiy) Khrebet, Koryakskiy Khrebet, Sredinnyy Khrebet, Verkhoyanskiy Khrebet, Momskiy Khrebet, Khrebet Suntar Khayata, Khrebet Cherskogo, Poluostrov Taygonos
67	Black Sea, S-Coast	Kuzey Anadolu Daglari / Pontus Mountains, Lesser Caucasus, Greater Caucasus
68	Caspian Sea, S-W-Coast	Agri Dagi, Süphan Dagi, Küh-e haye Sabalan, Alborz Mountains, Zagros Mountains, Lesser Caucasus, Greater Caucasus
69	Tigris-Euphrates	Mercan Daglari, Hakkari Daglari, Süphan Dagi, Zagros Mountains
70	Persian Gulf Coast	Zagros Mountains
71	Central Iran	Alborz Mountains, Zagros Mountains
72	Helmand	Hindu Kush
73	Farahrud	Hindu Kush
74	Caspian Sea, E-Coast	Hindu Kush, Alborz Mountains
75	Amu Darya	Zeravshan, Pamir-Alay, Turkestanskiy Khrebet, Gory Baysun Tau, Pamir, Karakorum, Hindu Kush, Malakand Range
76	Syr Darya	Turkestanskiy Khrebet, Pamir-Alai, Ferganskiy Khrebet, Chatkal'skiy Khrebet, Talas Alatau, Kyrgyz Ala Too, Terskey Ala Too, Kungey Ala Too,
77	Lake Balkash	Kyrgyz Ala Too, Kungey Ala Too, Ile Alatau, Borohoro-Shan, Dzhungarskiy Alatau, Narat Shan, Tien Shan
78	New Zealand	Ruapehu, Rolleston Range, Two Thumb Range, Liebig Range, Ben Ohau Range, Young Range, Olivine Range, Humboldt Mountains, Richardson Mountains, Livingstone Mountains

Indicator	Symbol	Input	Equation	Reference
Precipitation contribution WTU/basin	$P_T$	Average annual WTU precipitation sum (2001-2017): $P_{WTU}$ (km <sup>3</sup> )	$P_T = P_{WTU} / P_{BAS}$	32
		Average annual basin precipitation sum (2001-2017): $P_{BAS}$ (km <sup>3</sup> )		32
Inter-annual variability in precipitation	$P_{YV}$	Annual WTU precipitation for individual years (2001-2017): $P_Y$ (km <sup>3</sup> )	$P_{YV} = 1 - ((\max(P_Y) - \min(P_Y)) / \max(P_Y))$	32
Intra-annual variability in precipitation	$P_{MV}$	Average monthly WTU precipitation sum (2001-2017): $P_m$ (km <sup>3</sup> )	$P_{MV} = 1 - ((\max(P_m) - \min(P_m)) / \max(P_m))$	32
Precipitation	$P$	-	$P = 0.5 * (P_{YV} + P_{MV}) * P_T$	32
WTU snow cover	$S_T$	Average annual WTU snow cover: $S$ (-)		5
Inter-annual variability in snow cover	$S_{YV}$	Annual average WTU snow cover (2001-2017): $S_Y$ (-)	$S_{YV} = 1 - ((\max(S_Y) - \min(S_Y)) / \max(S_Y))$	5
Intra-annual variability in snow cover	$S_{MV}$	Average monthly snow cover (2001-2017): $S_m$ (-)	$S_{MV} = 1 - ((\max(S_m) - \min(S_m)) / \max(S_m))$	5
Snow	$S$	-	$S = 0.5 * (S_{YV} + S_{MV}) * S_T$	5
Glacier ice storage	$G_S$	Total glacier ice volume in WTU: $G_V$ (km <sup>3</sup> )	$G_S = G_V / (G_V + P_{WTU})$	48
		Average annual WTU precipitation sum (2001-2017): $P_{WTU}$ (km <sup>3</sup> )		32
Glacier melt yield	$G_M$	Average annual WTU precipitation sum (2001-2017): $P_{WTU}$ (km <sup>3</sup> )	$G_M = (P_{GLAC} - B) / (P_{GLAC} - B + P_{WTU})$	32
		Average annual precipitation sum glaciated area (2001-2017): $P_{GLAC}$ (km <sup>3</sup> )		32
		WTU average annual glacier mass balance: $B$ (km <sup>3</sup> )		17
				48
Glaciers	$G$	-	$G = G_S + G_M$	48
Lake and reservoir storage	$L$	Total volume stored in lakes and reservoirs in WTUs: $S_L$ (km <sup>3</sup> )	$L = S_L / (S_L + P_{WTU})$	50
		Average annual WTU precipitation sum (2001-2017): $P_{WTU}$ (km <sup>3</sup> )		32
Final supply index	$SI$		$(P + S + G + L) / 4$	


Indicator	Symbol	Input	Equation	Reference
Domestic demand	D <sub>DOM</sub>	Average annual downstream domestic water use (2001-2014): D <sub>DOM,y</sub> (km <sup>3</sup> )  Average monthly downstream domestic water use (2001-2014): D <sub>DOM,m</sub> (km <sup>3</sup> )  Average monthly P-ET (2001-2017) for downstream cells with domestic demand above threshold: WA <sub>DOM,m</sub> (km <sup>3</sup> ) Threshold is 1*10 <sup>-6</sup> km <sup>3</sup> per 0.05° grid cell	$\Sigma (D_{DOM,m} - WA_{DOM,m}) / D_{DOM,y}$	64
Industrial demand	D <sub>IND</sub>	Average annual downstream industrial water use (2001-2014): D <sub>IND,y</sub> (km <sup>3</sup> )  Average monthly downstream industrial water use (2001-2014): D <sub>IND,m</sub> (km <sup>3</sup> )  Average monthly P-ET (2001-2017) for downstream cells with industrial demand above threshold: WA <sub>IND,m</sub> (km <sup>3</sup> ) Threshold is 1*10 <sup>-6</sup> km <sup>3</sup> per 0.05° grid cell	$\Sigma (D_{IND,m} - WA_{IND,m}) / D_{IND,y}$	64
Irrigation demand	D <sub>IRR</sub>	Average annual downstream irrigation water use (2001-2014): D <sub>IRR,y</sub> (km <sup>3</sup> )  Average monthly downstream irrigation water use (2001-2014): D <sub>IRR,m</sub> (km <sup>3</sup> )  Average monthly P-ET (2001-2017) for downstream cells with irrigation demand above threshold: WA <sub>IRR,m</sub> (km <sup>3</sup> ) Threshold is 1*10 <sup>-6</sup> km <sup>3</sup> per 0.05° grid cell	$\Sigma (D_{IRR,m} - WA_{IRR,m}) / D_{IRR,y}$	64
Natural demand	D <sub>NAT</sub>	Average annual Environmental Flow Requirement at river basin outlet (2001-2014): D <sub>NAT,y</sub> (km <sup>3</sup> )  Average monthly Environmental Flow Requirement at river basin outlet (2001-2014): D <sub>NAT,m</sub> (km <sup>3</sup> )  Average monthly P-ET for downstream basin (2001-2017): WA <sub>NAT,m</sub> (km <sup>3</sup> )	$\Sigma (D_{NAT,m} - WA_{NAT,m}) / D_{NAT,y}$	51,52,64
Final demand index	DI		$(D_{IRR} + D_{IND} + D_{DOM} + D_{NAT}) / 4$	

640

641




642 **Extended Data Figures**




Extended Data Figure 1: a. Concept and global spread of Water Tower Units. The Water Tower Units (WTU) are defined as the intersection of Earth's major hydrological basins<sup>3</sup> and mountain ranges<sup>4</sup> meeting predefined thresholds for ice volume/area and or snow persistence (see Methods section). One WTU can consist of (parts of) multiple mountain ranges and one mountain range can be part of multiple WTUs. The example shows two hydrological basins in North America; Great Basin (red outline) and California (blue outline). The striped areas indicate two mountain ranges; the Sierra Nevada and the Cascade Range. The intersection of the hydrological basins and the mountain ranges defines the WTUs (dark tones). E.g. the Great Basin WTU is defined as the portion of the Sierra Nevada which is part of the Great Basin hydrological basin (dark red), and the California WTU is defined as the portion of the Sierra Nevada which is part of the California hydrological basin as well as a portion of the Cascade Range which is part of the California hydrological basin (dark blue). The WTU's dependent area (light tones) is defined as the sub-basins within the hydrological basin that are overlapping the WTU or downstream of sub-basins overlapping the WTU. b-e. The WTUs (dark tones) and associated WTU basins (light tones) for all 78 WTUs and WTU basins, grouped by continents: North America (a), Europe (b), Asia and Oceania (c), South America (d). Labels indicate the WTU IDs (see Extended Data Table 1 and 2 for corresponding names).

658




659




660

661 Extended Data Figure 2: Supply and Demand Index. a. The WTU supply index (blue colour ramp) and downstream  
662 demand index (brown colour ramp) for all 78 WTUs and WTU basins. b. Supply index (SI) and demand index (DI) for each  
663 WTU grouped per continent. Background color gradient indicates water tower importance (i.e. darker tones represent  
664 high SI and DI values). Points are labelled with WTU IDs (Extended Data Table 1 and 2, Extended Data Figure 1).

665




666




667

668 **Extended Data Figure 3: Annual precipitation and snow cover. a. Average annual precipitation between 2001 and 2017,**  
 669 **resampled bilinearly to 0.05° resolution based on ERA5<sup>32</sup>. b. Average snow persistence between 2001 and 2017,**  
 670 **resampled to 0.05° resolution based on MODIS MOD10CM1<sup>5</sup>.**

671



672




673


674 Extended Data Figure 4: Glacier ice and lake and reservoir volume. a. Total aggregated glacier ice volume, per WTU<sup>48</sup>. b.  
 675 Total aggregated lake and reservoir water volume per WTU<sup>50</sup>.

676






677




678

679 Extended Data Figure 5: Water use for irrigation and industry. a. Average annual irrigation water use per 0.05x0.05  
680 degrees grid cell 2001-2014<sup>73</sup>. b. Average annual industrial water use per 0.05x0.05 degrees grid cell 2001-2014<sup>73</sup>




681



682

683 Extended Data Figure 6: Domestic water use and natural water demand. a. Average annual domestic water use per  
 684  $0.05 \times 0.05$  degrees grid cell 2001-2014<sup>73</sup>. b. Total aggregated average annual natural water demand 2001-2014 per WTU  
 685 basin based on the Environmental Flow Requirement<sup>51,52,73</sup>

686



687

688 **Extended Data Figure 7: Sensitivity of WTU ranking to uncertainty in input data and indicator weights. Position change in**  
 689 **ranking of WTUs by WTI resulting from uncertainty in input data (blue), expressed as percentage of 1000 realizations of**  
 690 **the WTI index calculation. Position change in ranking of WTUs by WTI resulting from uncertainty in the weights of**  
 691 **individual indicators (red), expressed as percentage of 10,000 realizations of the WTI index calculation.**



## Supplementary Material

Supplementary Tables are provided as .xlsx spreadsheets

**Table S1: Indicator and index values per WTU. Overview of all supply indicators, supply index, demand indicators, demand index, and Water Tower Index. Explanations of all variables are provided in the Methods section and**

**Extended Data Table 3 and 4.**

**Table S2: Vulnerability and future change indicators per WTU. Overview of all vulnerability and future change indicators.**

**Explanations of all variables are provided in the Methods section.**